







AD-A185 053



THE GEORGE WASHINGTON UNIVERSITY

STUDENTS FACULTY STUDY R
ESEARCH DEVELOPMENT FUT
URE CAREER CREATIVITY CC
MMUNITY LEADERSHIP TECH
NOLOGY FRONTIF
ENGINEERING APP
ENC
GEORGE WASHIN
DTIC
SEP 2 3 1987



SCHOOL OF ENGINEERING AND APPLIED SCIENCE

87 9 13 039



EXPLICIT FORMULAE FOR THE DISTRIBUTIONS OF STOPPING VARIABLES UNDER WALD'S TRUNCATED SPRT FOR A POISSON PROCESS

bу

S. Zacks

GWU/IMSE/Serial T-516/87 6 May 1987

THE GEORGE WASHINGTON UNIVERSITY.
School of Engineering and Applied Science
Washington, DC 20052

Institute for Management Science and Engineering



Research Supported by Contract N00014-83-K-0216 Project NR 347 131 Office of Naval Research

This document has been approved for public sale and release; its distribution is unlimited.

SECURITY CLASSIFICATION OF THIS PAGE (When Date Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM		
1. REPORT NUMBER		3. RECIPIENT'S CATALOG NUMBER		
GWU/IMSE/Serial T-516/87	AD-A1850	153		
4. TITLE (and Subtitle)		S. TYPE OF REPORT & PERIOD COVERED		
EXPLICIT FORMULAE FOR THE DISTRI STOPPING VARIABLES UNDER WALD'S		SCIENTIFIC		
FOR A POISSON PROCESS	INUNCATED SERI	6. PERFORMING ORG. REPORT NUMBER		
100 11 1012500 1000255				
7. AUTHOR(a)		B. CONTRACT OR GRANT NUMBER(*)		
S. Zacks		N00014-83-K-0216		
	1			
9. PERFORMING ORGANIZATION NAME AND ADDRESS		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS		
GEORGE WASHINGTON UNIVERSITY				
INSTITUTE FOR MANAGEMENT SCIENCE	AND ENGINEERING			
WASHINGTON, DC 20052		12. REPORT DATE		
OFFICE OF NAVAL RESEARCH		6 May 1987		
CODE 411 S&P ARLINGTON, VA 22217		13. NUMBER OF PAGES		
14. MONITORING AGENCY NAME & ADDRESS(II dillerent	from Controlline Office)	17		
14. MONITORINO AGENCY HAME & ADDRESS, MINORIN				
•		UNCLASSIFIED		
		18a. DECLASSIFICATION/DOWNGRADING SCHEDULE		
16. DISTRIBUTION STATEMENT (of this Report)				
APPROVED FOR PUBLIC SALE AND RE	LEASE; DISTRIBUT	CION UNLIMITED.		
17. DISTRIBUTION STATEMENT (of the abstract entered in	in Block 20, if different fre	m Report)		
•				
18. SUPPLEMENTARY NOTES				
19. KEY WORDS (Continue on reverse side if necessary and	d identify by block number)	'		
STOPPING VARIABLES	POISSON PROCESS	SES		
WALD SPRT	MTBF			
		1		
20 ACCENCE (Continue or annual day)	I Identify by black symbol			
20. ABSTRACT (Continue on reverse elde il necessary and Explicit formulae are devel				
variables associated with truncated versions of the Wald sequential prob-				
ability ratio tests (SPRT) of hy	-			
(MTBF) of a Poisson process. The expressions for linear boundarie				
		also be applied for determi-		
ning confidence intervals for th	e MTBF after seq	quential stopping.		

THE GEORGE WASHINGTON UNIVERSITY School of Engineering and Applied Science Washington, DC 20052

Institute for Management Science and Engineering

EXPLICIT FORMULAE FOR THE DISTRIBUTIONS
OF STOPPING VARIABLES UNDER WALD'S TRUNCATED SPRT
FOR A POISSON PROCESS

bу


S. Zacks

Abstract of GWU/IMSE/Serial T-516/87 6 May 1987

Explicit formulae are developed for the distributions of the stopping variables associated with truncated versions of the Wald sequential probability ratio tests (SPRT) of hypotheses about the mean time between failures (MTBF) of a Poisson process. These formulae are based on newly derived expressions for linear boundaries crossing probabilities under Poisson processes. The results of the present study can also be applied for determining confidence intervals for the MTBF after sequential stopping.

Key Words: Stopping variables, Wald SPRT, Poisson processes, MTBF

Research Supported by Contract N00014-83-K-0216 Project NR 347 131 Office of Naval Research



THE GEORGE WASHINGTON UNIVERSITY School of Engineering and Applied Science Washington, DC 20052

Institute for Management Science and Engineering

EXPLICIT FORMULAE FOR THE DISTRIBUTIONS
OF STOPPING VARIABLES UNDER WALD'S TRUNCATED SPRT
FOR A POISSON PROCESS

bу

S. Zacks

GWU/IMSE/Serial T-516/87 5 May 1987 SSSS CONTROL AND PROPERTY OF THE PROPERTY OF T

1. Introduction

The computation of probabilities associated with sequential stopping variables can often be easily performed by recursive techniques (e.g. Armitage et al (1969), Aroian and Robinson (1969), Samuel-Cahn (1974), Pocock (1977), Zacks (1974, 1980, 1981, 1987)). Explicit formulae for probabilities of boundary crossings are generally difficult to obtain. Wiener process and other asymptotic approximations to the distributions of stopping times are discussed by Siegmund (1985). The present paper shows that in the case of sequential procedures based on Poisson processes, explicit expression can be derived. These derivations are facilitated by the nondecreasing nature of the sample paths, the memory-less property of the Poisson process, and the possibility to solve certain recursive relations associated with the probability distributions of the stopping variables. For the purpose of showing the potential of this technique, we derive explicit expressions for the cumulative distribution functions (CDF) of stopping variables associated with the acceptance and with the rejection boundaries of a Wald's SPRT, which is frequency truncated. Such stopping boundaries are applied in reliability life testing for systems having

exponentially distributed life length (see MIL-STD 781C (1977)). Methods of determining confidence intervals for the MTBF, after such sequential life testing, were given by Siegmund (1978), Bryant and Schmee (1979), and Zacks (1987). Siegmund approximated the distributions of stopping times, while Zacks determined them recursively. The results of the present paper eliminate the need for recursive computations in the Poisson case.

In Section 2 we present the stopping boundaries, the observed process and some general definitions. In Section 3 we derive explicitly certain boundary crossing probabilities. Section 4 is devoted to an actual derivation of the distributions of the stopping variables, for a particular case under consideration. A numerical example is provided in Section 5.

2. The Failure Process and Stopping Variables

Let $\{X(t): \theta < t < \infty\}$ be a Poisson process with intensity λ , $0 < \lambda < \infty$; i.e., $E_{\lambda}(X(t)) = \lambda t$, $0 < t < \infty$. We consider a truncated Wald SPRT, which is specified by the boundary lines

(2.1)
$$b_{R}(t) = \begin{cases} k_{1} + bt, & 0 < t \le t_{k_{g}-k_{1}} \\ k_{g}, & t_{k_{g}-k_{1}} \le t \le t_{k_{g}+k_{0}} \end{cases}$$

and

(2.2)
$$b_A(t) = -k_0 + bt, \quad t_{k_0} \le t \le t_{k_0+k_s}$$

where k_1 , k_0 , k_s are integers, $k_s > k_1$, and $t_1 = i/b$ (i = 1, 2, ...). The boundary $b_A(t)$ is called the acceptance boundary, and $b_R(t)$ is called the rejection boundary. Notice that $b_A(t)$ and $b_R(t)$ intersect at $(t_{k_0+k_s}, k_s)$.

We define the stopping times (variables)

(2.3)
$$\tau_{R} = \inf \{t: X(t) \ge b_{R}(t), 0 < t < t_{k_0 + k_s} \},$$

(2.4)
$$\tau_{A} = \inf \{t: X(t) = b_{A}(t), t_{k_{0}} \leq t \leq t_{k_{0}+k_{S}} \},$$

where $\{X(t): t > 0\}$ is a nondecreasing jump process, with jumps of one unit. The process is stopped at the first instant τ at which it crosses either the lower boundary, $b_A(t)$, or the upper boundary $b_R(t)$. Obviously, $\tau = \min\{\tau_A, \tau_R\}$.

For the derivation of the distributions of the stopping times τ_R , τ_A and τ one has to distinguish between three cases,

Case I:
$$k_1 < k_8 \le k_1 + k_0$$
;
Case II: $k_1 + k_0 < k_8 \le 2k_1 + k_0$;
Case III: $2k_1 + k_0 < k_8$.

Since the method of derivation of the distributions is similar in all the three cases, we focus attention in the present paper only on Case I.

The sample paths of the Poisson process $\{X(t): 0 < t\}$ assume only nonnegative integer values. Thus, a sample path can cross the lower boundary $b_A(t)$ only at the discrete times t_i , $k_0 \le i \le k_8 + k_0 - 1$. Let

$$f_{A,k_0}^{(\lambda)}(j) = P_{\lambda} \{ \tau_A - t_{k_0+j} \}$$

be the probability distribution function of the discrete stopping variable τ_A , over $t_{k_0+j^*}$. We introduce the notation

$$I\{A\} = \begin{cases} 1, & \text{if A is true} \\ 0, & \text{otherwise;} \end{cases}$$

and $p(j;\xi)$ and $Pos(j;\xi)$ are to be, respectively, the PDF and CDF of the Poisson distribution with mean ξ .

Let $F_A^{(\lambda)}(t)$ denote the corresponding CDF of τ_A . The stopping time τ_R on the other hand, has an absolutely continuous distribution. Define the probability function

(2.5)
$$g_{\lambda}(j; t) = P_{\lambda}\{X(t) = j, \tau \geq t\}, j = 0, 1,...$$

Let [a] be the largest integer smaller than or equal to a, and define

(2.6)
$$M(t) = \begin{cases} [k_1 + bt], & \text{if } k_1 + bt \text{ is not an integer} \\ k_1 + bt - 1, & \text{otherwise.} \end{cases}$$

Similarly, let $a^+ = max (a, 0)$ and let

(2.7)
$$m(t) = \begin{cases} [-k_0 + bt], & \text{if } t \ge t_{k_0} \\ [-k_0 + bt]^+ - 1, & \text{if } t < t_{k_0} \end{cases}$$

Then

(2.8)
$$P_{\lambda}\{\tau > t\} = \sum_{j=m(t)+1}^{M(t)} g_{\lambda}(j; t).$$

Let $F_R^{(\lambda)}$ designate the CDF of τ_R , then

(2.9)
$$F_{R}^{(\lambda)}(t) = 1 - F_{A}^{(\lambda)}(t) - P_{\lambda}\{\tau > t\}, \quad 0 < t.$$

Furthermore, from the Markovian property of the Poisson process, for each $t_{i-1} < t \le t_i$

(2.10)
$$F_R^{(\lambda)}$$
 (t) = $F_R^{(\lambda)}$ $\{t_{i-1}\}$

$$+\sum_{j=m(t_{i-1})+1}^{M(t_{i-1})} g_{\lambda}(j; t_{i-1}) \left[1 - Pos(M(t_{i-1}) + 1 - j; \lambda(t - t_{i-1}))\right]$$

This formula holds for i=1 too, if we define $g_{\lambda}(j;\,t_0)=I\{j=0\}$. Notice that

$$F_R^{(\lambda)}$$
 (t) is a differentiable function of t, for $t_{i-1} < t < t_i$ (i = 1, 2,..., $k_s + k_0$).

Thus, $F_R^{(\lambda)}(t)$ is absolutely continuous. In Section 4 we develop explicit formulas for $F_A^{(\lambda)}(t)$ and $F_D^{(\lambda)}(t)$, appropriate for Case I.

3. Linear Boundaries Crossing Probabilities

Let h_{λ} (i, s) designate the probability that a Poisson process, with intensity λ , assumes the value i at time s, and does not cross the linear boundary x(u) = bu for $0 < u \le s$, i. e.,

(3.1)
$$h_{\lambda}(i; s) = P_{\lambda} \{ \sup(X(u) - bu < 0; 0 < u < s), X(s) = i \}.$$

Let \mathfrak{A}_{s}^{-} be the region below x(u) over (0, s], and \mathfrak{A}_{s}^{+} the region above it (including the boundary), i. e.,

(3.2)
$$\mathbf{U}_{\mathbf{s}}^{-} = \{(\mathbf{u}, \mathbf{x}): 0 \le \mathbf{x} < \mathbf{b}\mathbf{u}, 0 < \mathbf{u} \le \mathbf{s}\}$$

and

(3.3)
$$u_{\mathbf{g}}^{+} = \{(\mathbf{u}, \mathbf{x}): \ \mathbf{b}\mathbf{u} \leq \mathbf{x}, \quad 0 < \mathbf{u} \leq \mathbf{s}\}.$$

Conditioning on the level at which the last entrance of a sample path to \mathbf{u}_{s}^{-} occurs, we obtain the recursive equation

$$(3.4) \quad h_{\lambda}(i;\,\mathbf{s}) = \left\{ \begin{array}{l} 0, & \text{if } i=0 \\ \\ 0, & \text{if } i>0, \, s\leq t_i \end{array} \right.$$

$$p(i;\,\lambda s) = \sum_{\ell=1}^{i} p(\ell;\,\lambda t_{\ell}) \; h_{\lambda}(i-\ell;\,s-t_{\ell}), \; \text{if } i>0, \, s>t_i$$
 Let $\mu=\lambda/b$ and let $H_{\lambda}(i;\,s) = \sum_{j=0}^{i} h_{\lambda}(j;\,s)$. Thus, for each $j=0,\,1,\,2,\ldots$

and i = j + 1, j + 2,... we obtain the recursive relation

(3.5)
$$H_{\lambda}(j; t_{i}) = \begin{cases} p(0; i\mu), & j = 0 \\ Pos(j; i\mu) - \sum_{\ell=1}^{j} p(\ell; \mu\ell) H_{\lambda}(j - \ell; t_{i-\ell}), j \geq 1 \end{cases}$$

An explicit solution of (3.5) can be obtained by using the theory of formal power-series (see P. Henrici (1974); p. 17).

For $\theta \in (-1, 1)$ and $\delta = 0, 1, 2,...$ define the power-series:

(3.6)
$$H_{\lambda,\delta}^{*}(\theta) = \sum_{j=0}^{\infty} H_{\lambda}(j; t_{j+\delta}) \theta^{j},$$

$$F_{\lambda,\delta}^{*}(\theta) = \sum_{j=0}^{\infty} Pos(j; \mu(j+\delta)) \theta^{j},$$

$$P_{\lambda,\delta}^{*}(\theta) = \sum_{j=0}^{\infty} p(j; \mu(j+\delta)) \theta^{j},$$

where $h_{\lambda}(0; 0) = 1$ and p(0, 0) = 1. The power-series $P_{\lambda,0}^{*}(\theta)$ has an inverse,

given by
$$Q_{\lambda}^{*}(\theta) = \sum_{n=0}^{\infty} q_{n}^{(\lambda)} \theta^{n}$$
, where

(3.7)
$$q_{n}^{(\lambda)} = \begin{cases} 1, & n = 0 \\ -p(1; \mu), & n = 1 \end{cases}$$
$$(-1)^{n} D_{n}(\mu), & n \geq 2$$

where $D_n(\mu)$ is the determinant of the n x n matrix

(3.8)
$$A_{\Pi}(\mu) = \begin{bmatrix} p(1; \mu) & p(2; 2\mu) & \dots & p(n; n\mu) \\ 1 & p(1; \mu) & \dots & p(n-1; (n-1)\mu) \\ 0 & 1 & \dots & p(n-2; (n-2)\mu) \\ 0 & 0 & \dots & p(n-3; (n-3)\mu) \\ \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & p(1; \mu) \end{bmatrix}$$

From (3.5) we obtain the equation

(3.9)
$$H_{\lambda,\delta}^{*}(\theta) = F_{\lambda,\delta}^{*}(\theta) - H_{\lambda,\delta}^{*}(\theta) (P_{\lambda,0}^{*}(\theta) - 1),$$

or, equivalently,

(3.10)
$$H_{\lambda,\delta}^{*}(\theta) P_{\lambda,0}^{*}(\theta) = F_{\lambda,\delta}^{*}(\theta).$$

Hence,

(3.11)
$$H_{\lambda,\delta}^{*}(\theta) = F_{\lambda,\delta}^{*}(\theta) Q_{\lambda}^{*}(\theta)$$

and, for each i = 0, 1,...

(3.12)
$$H_{\lambda}(i; t_{i+\delta}) = \sum_{n=0}^{i} q_{n}^{(\lambda)} Pos(i-n; (i-n+\delta)\mu).$$

In a similar fashion we obtain that for every j = 0, ..., i - 1 (i = 1, 2, ...)

(3.13)
$$h_{\lambda}(j; t_{i}) = \sum_{n=0}^{j} q_{n}^{(\lambda)} p(j-n; (i-n)\mu).$$

Consider a linear boundary $x_{\ell}(u) = -\ell + bu$, for an integer value ℓ . Let $f_{A,\ell}^{(\lambda)}(i)$

be the probability that the process crosses this boundary for the first time at $(t_{\ell+i}, i)$, i = 0, 1, 2, ... These boundary crossing probabilities satisfy the following recursive equation for i = 0, 1, ...

(3.14)
$$f_{A,\ell}^{(\lambda)}(i) = p(i; (\ell + i)\mu) - I\{i > 1\} \sum_{j=0}^{i-1} f_{A,\ell}^{(\lambda)}(j) p(i - j; (i - j) \mu).$$

Define the power-series, for $\theta \in (-1, 1)$,

(3.15)
$$f_{\lambda,\ell}^{*}(\theta) = \sum_{i=0}^{\infty} f_{A,\ell}^{(\lambda)}(i) \theta^{i}.$$

The recursive equation (3.14) yields the equation

$$f_{\lambda,\ell}^{*}(\theta) = P_{\lambda,\ell}^{*}(\theta) \left[P_{\lambda,0}^{*}(\theta)\right]^{-1},$$

where $\left(P_{\lambda,0}^*(\theta)\right)^{-1} \equiv Q_{\lambda}^*(\theta)$. Hence, we obtain by multiplying the power-series,

(3.17)
$$f_{A,\ell}^{(\lambda)}(i) = \sum_{n=0}^{1} q_n^{(\lambda)} p(i-n; (\ell+i-n)\mu), \text{ for } i=0,1,$$

4. The Distributions of Stopping Times in Case I

The continuation region (between $b_A(t)$ and $b_R(t)$) is partitioned into four regions:

$$\begin{split} & R_1 = \big\{ (t,\,x) \colon \quad 0 < t \le t_{k_S-k_1}, \quad 0 \le x < b_R(t) \big\}; \\ & R_2 = \big\{ (t,\,x) \colon \ t_{k_S-k_1} < t \le t_{k_0}, \quad 0 \le x < b_R(t) \big\}; \\ & R_3 = \big\{ (t,\,x) \colon \ k_0 < t \le t_{k_0+k_1}, \quad b_A(t) < x < b_R(t) \big\}; \text{ and } \\ & R_4 = \big\{ (t,\,x) \colon \ t_{k_0+k_1} < t \le t_{k_S+k_0}, \ b_A(t) < x < b_R(t) \big\}. \end{split}$$

4.1 Distributions over R₁

Over R_1 , $\tau=\tau_R$, while $P_{\lambda}\{\tau=\tau_A\}=0$. From the Markovian properties we obtain, for $i=1,...,k_8-k_1$, $j=0,1,...,k_1+i-1$,

(4.1)
$$g_{\lambda}(j; t_i) = p(j; \mu i)$$

$$- I\{j \geq k_1 + 1\} \sum_{\ell=1}^{j-k_1} p(k_1 + \ell; \ell \mu) h_{\lambda}(j - k_1 - \ell; t_i - t_{\ell}).$$

Hence, for $1 \le i \le k_S - k_1$,

$$\begin{array}{ll} \text{(4.2)} & P_{\lambda} \big\{ \tau_k > t_i \big\} & = \text{Pos}(k_1 + i - 1; \, i\mu) \\ \\ & - I \big\{ i \geq 2 \big\} \displaystyle \sum_{j=k_1+1}^{k_1+i-1} \sum_{\ell=1}^{j-k_1} p(k_1 + \ell; \, \ell\mu) \, \, h_{\lambda}(j-k_1 - \ell; \, t_i - t_{\ell}). \end{array}$$

Changing the order of summation we obtain

$$(4.3) P_{\lambda} \{ \tau_{R} > t_{1} \} = Pos(k_{1} + i - 1; i\mu)$$

$$- I\{i \geq 2\} \sum_{\ell=1}^{i-1} p(k_{1} + \ell; \ell\mu) H_{\lambda}(i - 1 - \ell; t_{1-\ell}).$$

Thus, for every $i = 1,..., k_S - k_1$, we obtain from (4.3) and (3.12),

$$(4.4) \quad F_{A}^{(\lambda)}(t_{i}) = 1 - Pos(k_{1} + i - 1; i\mu)$$

$$+ I\{i \geq 2\}$$

$$\cdot \sum_{\ell=1}^{i-1} p(k_{1} + \ell; \ell\mu) \sum_{n=0}^{i-1-\ell} q_{n}^{(\lambda)} \quad Pos(i - 1 - \ell - n; (i - \ell - n)\mu).$$

4.2 The Distributions over R2

Over $\rm R_2$ we obtain similar results to those over $\rm R_1$, with only slight changes. Here, the CDF of $\tau_{\rm A}$ is

(4.5)
$$F_{A}^{\lambda}(t_{i}) = \begin{cases} 0, & i = k_{s} - k_{1} + 1, ..., k_{0} - 1 \\ f_{A,k_{0}}^{(\lambda)} & (0), & i = k_{0}. \end{cases}$$

The CDF of au_R is given by

(4.6)
$$F_{R}^{(\lambda)}(t_{i}) = 1 - Pos(k_{s} - 1; i\mu)$$

 $+ I\{k_{s} > k_{1} + 1\} \sum_{\ell=1}^{k_{s}-k_{1}-1} p(k_{1} + \ell; \ell\mu) H_{\lambda}(k_{s} - k_{1} - \ell - 1; t_{i-\ell})$

for $i = k_8 - k_1 + 1,..., k_0$. According to (3.12),

(4.7)
$$H_{\lambda}(k_{S} - k_{1} - \ell - 1; t_{i-\ell})$$

$$= \sum_{n=0}^{k_{S}-k_{1}-\ell-1} q_{n}^{(\lambda)} \operatorname{Pos}(k_{S} - k_{1} - \ell - n - 1; (i - \ell - n)\mu).$$

4.3 The Distributions over R3

The probability distribution (PDF) of $\tau_{\rm A}$ over R₃ is given by

(4.8)
$$P_{\lambda} \{ \tau_{A} = k_{0} + i \} = f_{A,k_{0}}^{(\lambda)}(i), i = 1,..., k_{1},$$

where $f_{A_sk_0}^{(\lambda)}$ (i) is given by (3.17).

The probabilities $g_{\lambda}(j; t_{k_0+i})$, for $i \le i \le k_1$, are given by

$$(4.9) \quad \mathbf{g}_{\lambda}[\mathbf{j}; \, \mathbf{t}_{\mathbf{k}_{0}} + \mathbf{i}] = \mathbf{I}\{1 \leq \mathbf{j} \leq \mathbf{k}_{1}\}$$

$$\cdot \left[\mathbf{p}[\mathbf{j}; \, \mu(\mathbf{k}_{0} + \mathbf{i})] - \sum_{\ell=0}^{i-1} f_{\mathbf{A}, \mathbf{k}_{0}}^{(\lambda)}(\ell) \, \mathbf{p}[\mathbf{j} - \ell; \, (\mathbf{j} - \ell)\mu] \right]$$

$$+ \mathbf{I}\{\mathbf{k}_{1} + 1 \leq \mathbf{j} \leq \mathbf{k}_{1} - 1\} \cdot \mathbf{p}[\mathbf{j}; \, \mu(\mathbf{k}_{0} + \mathbf{i})]$$

$$- \mathbf{I}\{\mathbf{k}_{1} + 1 \leq \mathbf{j} \leq \mathbf{k}_{1} - 1\}$$

$$\cdot \sum_{\ell=0}^{i-1} f_{\mathbf{A}, \mathbf{k}_{0}}^{(\lambda)}(\ell) \, \mathbf{p}[\mathbf{j} - \ell; \, \mu(\mathbf{i} - \ell)]$$

$$- \mathbf{I}\{\mathbf{k}_{1} + 1 \leq \mathbf{j} \leq \mathbf{k}_{1} - 1\}$$

$$\cdot \sum_{m=1}^{j-k} \mathbf{p}[\mathbf{k}_{1} + m; \, m\mu] \, \mathbf{h}_{\lambda}[\mathbf{j} - \mathbf{k}_{1} - m; \, \mathbf{t}_{\mathbf{k}_{0} + \mathbf{i} - m}]$$

where, according to (3.13), for $k_1 + 1 \le j \le k_s - 1$,

(4.10)
$$h_{\lambda}[j-k_{1}-m; t_{k_{0}+i-m}]$$

$$= \sum_{n=0}^{j-k_{1}-m} q_{n}^{(\lambda)} p[j-k_{1}-m-n; \mu(k_{0}+i-m-n)].$$

From (4.9) we obtain, for $i \le i \le k_1$,

$$\begin{aligned} \text{(4.11)} \quad & F_{R}^{(\lambda)} \ (t_{i}) = 1 - F_{A}^{(\lambda)} \ (t_{i}) - P_{\lambda} \big\{ \tau > t_{i} \big\} \\ & = 1 - F_{A}^{(\lambda)} \ (t_{i}) - Pos(k_{S} - 1; \ \mu(k_{0} + i)) + Pos(i; \ \mu(k_{0} + i)) \\ & + \sum_{\ell=0}^{i-1} f_{A,k_{0}}^{(\lambda)} (\ell) \ [Pos(k_{S} - 1 - \ell; \ \mu(i - \ell)) - Pos(i - \ell; \ \mu(i - \ell))] \\ & + I \big\{ k_{S} > k_{1} + 1 \big\} \\ & \cdot \sum_{\ell=1}^{k_{S}-k_{1}-1} p(k_{1} + \ell; \ \ell\mu) \ H_{\lambda} \big[k_{S} - k_{1} - \ell - 1; \ t_{k_{0}+i-\ell} \big]. \end{aligned}$$

4.4 The Distributions over R4

We assume that $k_S>k_1+1$. If $k_S=k_1+1$ then $P_\lambda \big\{ \tau \leq t_{k_0+k_1} \big\}=1$.

Applying the Markovian property of the Poisson process, we obtain for

$$1 \leq i \leq k_8 - k_1 - 1,$$

(4.12)
$$P_{\lambda} \{ \tau_{A} = t_{k_{0}+k_{1}} + i \} = f_{A,k_{0}}^{(\lambda)} (k_{1} + i)$$

$$= \sum_{\ell=1}^{i} g_{\lambda} [k_{1} + \ell; t_{k_{0}+k_{1}}]$$

$$\cdot \left\{ p(i - \ell; \mu_{i}) - 1 \{i > \ell\} \sum_{r=0}^{i-\ell-1} f_{A,\ell}^{(\lambda)} (m) p[i - \ell - m; \mu(i - \ell - m)] \right\}$$

where $g_{\lambda}(k_1 + \ell; t_{k_0+k_1})$ is given by (4.9).

In a similar manner we establish that, for every $1 \le i \le k_S - k_1 - 2$,

$$(4.13) \ \mathbf{g}_{\lambda}[k_{1} + j; \, t_{k_{0}+k_{1}+i}] = I\{i + 1 \leq j \leq k_{s} - k_{1} - 1\} \sum_{\ell=1}^{j} \mathbf{g}_{\lambda}[k_{1} + \ell; \, t_{k_{0}+k_{1}}]$$

$$\cdot \left[p(j - \ell; \, \mu_{i}) - I\{\ell < i\} \right] \sum_{k=1}^{j} f_{\mathbf{A},\ell}^{(\lambda)} (m) \, p[j - \ell - m; \, \mu(i - \ell - m)] .$$

Hence.

$$(4.14) P_{\lambda} \{ \tau > t_{k_0 + k_1 + i} \} = I \{ i < k_8 - k_1 - 1 \} \sum_{j=i+1}^{k_8 - k_1 - 1} g_{\lambda} [k_1 + j; t_{k_0 + k_1 + i}]$$

$$= I \{ i < k_8 - k_1 - 1 \}$$

$$\left\{ \sum_{\ell=1}^{j} g_{\lambda} \left[k_{1} + \ell; t_{k_{0}+k_{1}} \right] \left[Pos \left(k_{S} - k_{1} - \ell - 1; i\mu \right) - Pos (i - \ell; i\mu) \right] \right.$$

$$\left. + \sum_{\ell=i+1}^{k_{S}-k_{1}-1} g_{\lambda} \left[k_{1} + \ell; t_{k_{0}+k_{1}} \right] Pos \left(k_{S} - k_{1} - \ell - 1; i\mu \right) \right.$$

$$- \ I\{i > 1\} \sum_{m=0}^{i-2} \sum_{\ell=1}^{i-1-m} \ g_{\lambda}[k_1 + \ell; \ t_{k_0+k_1}] \cdot f_{A,\ell}^{(\lambda)}(m)$$

$$-\left[\operatorname{Pos}(k_{S}-k_{1}-\ell-m-1;\,\mu(i-\ell-m))-\operatorname{Pos}(i-\ell-m;\,\mu(i-\ell-m))\right].$$

SASASON/ POCCOCCO ESSASAS POCCOCCO POCCOCO POCCOCO POCCOCO POCCOCO POCCOCO POCCOCO POCCOCO POCOCO POCOCOCO POCOCOCO

5. Numerical Example

It is a simple matter to compute the CDF of the stopping times (variables) τ_R , τ_A and τ according to the formula given in Section 4. In Table 5.1 we present these distributions for Case I, with $k_0 = 15$, $k_1 = 15$, $k_2 = 28$ and $k_3 = 20$. We present $F_R^{(\lambda)}(t)$, $F_A^{(\lambda)}(t)$ and $F_R^{(\lambda)}(t)$ at the values of $k_1 = k_2 = 10$. Notice (i = 1, 2, ..., $k_2 + k_3 = 10$) for Poisson processes with $k_3 = 10$ and $k_3 = 10$. Notice that the values of $k_3 = 10$ are specified for a convenient time unit, which is traditionally take to be the MTBF under the alternative hypothesis, and a specified number of systems on test (see Zacks (1987)). The values of Table 5.1 were computed on a PC, using a TURBO-PASCAL program, which is available upon request.

COLOCOCO EXPERSO ESCOCIOS PARAMENTOS ESCOSOS

Table 5.1: Cumulative Distributions of Stopping Times For Truncated Wald SPRT $(k_0=k_1=15, k_S=28, B=20)$

		λ = 10		λ = 20		
t	F _R (t)	F _A (t)	F(t)	F _R (t)	F _A (t)	F(t)
Ø.Ø5	0.000000	0.000000	0.000000	0.000000	0.000000	0.000000
0.10	0.000000	0.00 0000	0.000000	0.000000	0.000000	0.000000
Ø.15	0.000000	0. 000000	0.000000	Ø.ØØØØØØ	0.000000	0.000000
0.20	0.000000	0.000000	0.000000	0.000000	0.000000	0.000000
Ø.25	0.000000	0.000000	0.000000	0.000000	0.000000	0.000000
Ø.3Ø	0.000000	0.000000	0.000000	0.000002	0.000000	0.000002
Ø.35	0.000000	0.000000	0.000000	0.000005	0.000000	0.000005
0.40	0.000000	0.000000	0.000000	0.000013	0.000000	0.000013
Ø.45	0.000000	0.000000	0.000000	0.000028	0.000000	0.000028
0.50	0.000000	6.6 66666	0.000000	0.000056	0.000000	0.000056
Ø.55	0.000000	0.000000	0.000000	0.000100	0.000000	0.000100
0.60	0.000000	0.000000	0.000000	0.000165	6.000000	0.000165
Ø.65	0.000000	0.000000	0.000000	0.000258	0.000000	0.000258
Ø.70	0.000000	0.000000	0.000000	0.000661	6.000000	0.000661
Ø.75	0.000000	Ø.ØØØ553	0.000553	0.001728	0.000000	0.001729
Ø.8Ø Ø.85	0.000000	Ø.ØØ3Ø69	0.003069	0.004111 0.008836	0. 000002 0. 000007	0.004 113 0.008844
Ø.9Ø	0.000000	0.009555 0.022050	0.009555	0.017319	0.000007 0.000020	Ø.Ø17339
Ø.95	0.000001	Ø.Ø42105	0.022050 0.042106	Ø.Ø31269	0.000020 0.000044	0.031312
1.00	0.000001	Ø.070480	Ø.Ø7Ø482	Ø.Ø52481	Ø.000044 Ø.000085	Ø.Ø52566
1.05	0.000005	Ø.107088	Ø.107094	0.082541	0.000149	Ø.Ø8269Ø
1.10	0.000013	Ø.151118	Ø.151131	0.122503	0.000243	Ø.122747
1.15	0.000027	0.201242	Ø.2Ø1269	Ø.172631	Ø.ØØØ373	0.173004
1.20	0.000056	Ø.255844	0.255900	Ø.232258	0.000545	0.232803
1.25	0.000109	Ø.31323Ø	0.313340	Ø.299814	0.000764	0.300578
1.30	0.000204	Ø.371778	0.371983	Ø.372996	0.001035	Ø.374Ø31
1.35	0.000367	0.430045	0.430412	0.449057	0.001362	0.450420
1.40	0.000635	Ø.486825	Ø.48746Ø	Ø.525136	0.001749	Ø.526885
1.45	0.001061	Ø.541171	Ø.542232	0.598567	0.0 02198	0.600766
1.50	0.001716	Ø.592389	0.594105	Ø.667131	0.002712	Ø.669843
1.55	0.002692	Ø.64ØØ16	0.642707	Ø.729195	0.003291	0.732486
1.60	0.004105	Ø.683788	0. 687893	Ø.783771	0.003936	0.787707
1.65	0.015958	0.713544	0.729502	0.830488	0.004514	0.835003
1.70	0.029219	0.737448	Ø.766667	Ø.869395	0.005117	0.874512
1.75	0.042745	Ø.758349	0.801093	0.901008	0.005788	0.906796
1.80	0.056490	Ø.777129	Ø.833618	0.926085	0.006545	0.932630
1.85	0.070690	0.794002	0.864692	0.945507	0.007393	0.952899
1.90	0.085617	0.809013	Ø.894631	0.960181	0.008329	Ø.96851Ø
1.95	0.101469	Ø.822211	0.923680	0.970978	0.009350	0.980328
2.00	0.118164	Ø.833688	Ø.951852	0.978670	0.010451	Ø.989121
2.05	0.134752	Ø.843573	0.978325	0.983895	0.011627	0.995522
2.10	0.147980	0.852020	1.000000	0.987127	0.012873	1.000000

PERCESSION NAMED IN TAXABLE

essential proposorial more second mesos resultante de la constante de la const

REFERENCES

- [1] Armitage, P., C. K. McPherson and B. C. Rowe (1969). Repeated significance tests on accumulating data. J. Roy. Statist. Soc. Ser. A: 235-244.
- [2] Aroian, L. A. and D. E. Robinson (1969). Direct methods for exact truncated sequential tests of the mean of a normal distribution. *Technometrics*, 11, 661-675.
- [3] Bryant, C. M. and J. Schmee (1979). Confidence limits on MTBF for sequential test plans of MIL-STD 781. Technometrics, 21: 33-42.
- [4] Henrici, P. (1974). Applied and Computational Complex Analysis, Vol. I. John Wiley, New York.
- [5] MIL-STD 781C (1977). Reliability Design Qualifications and Production Acceptance Tests: Exponential Distribution. Department of Defense, Washington DC.
- [6] Samuel-Chan, E. (1974). Repeated significance tests II, for hypotheses about the normal distribution, Commun. Statist. 3. 711-733.
- [7] Siegmund, D. (1978). Estimation Following Sequential Tests, Blometrika. 65, 341-349.
- [8] Siegmund, D. Sequential Analysis: Tests and Confidence Intervals. Springer-Verlag, New York, 1985.
- [9] Zacks, S. (1974). The proportional closeness and expected sample size of sequential procedures for estimating tail probabilities in exponential distributions, Commun. Statist. 3, 105-120.
- [10] Zacks, S. (1980). Distribution of stopping variables in sequential procedures for the detection of shifts in the distributions of discrete random variables. Commun. Statist. B1, 1-8.
- [11] Zacks, S. (1981). The probability distribution and expected value of a stopping variables associated with one-sided CUSUM procedures for nonnegative integer valued random variables. Commun. Statist.

 A10. 2245-2258.
- [12] Zacks, S. (1987). The determination of confidence intervals for the MTBF of systems having exponential life distribution after sequential testing. (Submitted for publication).

OFFICE OF NAVAL RESEARCH MATHEMATICAL SCIENCES DIVISION

BASIC DISTRIBUTION LIST FOR UNCLASSIFIED TECHNICAL REPORTS

OCTOBER 1985

copies	Copies
Mathematical Sciences	Navy Library
Division (Code 411)	National Space Technology Laboratory
Office of Naval Research	ATIN: Navy Librarian
Arlington, VA 22217-5000 3	Bay St. Louis, MS 39522 1
Defense Technical Information	U.S. Army Research Office
Center	P.O. Box 12211
Cameron Station Alexandria, VA 22314 12	ATIN: Dr. J. Chandra
·	Research Triangle Park, NC 22706 1
Commanding Officer	
Office of Naval Research	Director
Eastern/Central Regional Office	National Security Agency
ATIN: Director for Science	ATIN: R51, Dr. Maar
Barries Building	Fort Meade, MD 20755 1
495 Summer Street	ATTA A CT
Boston, MA 02210 1	ATAA-SL, Library
U.S. ONR Liaison Office - Far East	U.S. Army TRADOC Systems
ATIN: Scientific Officer	Analysis Activity
APO San Francisco 96503 1	Department of the Army
	White Sands Missile Range, NM 88002
Applied Mathematics Laboratory	
David Taylor Naval Ship Research	Library, Code 1424
and Development Center	Naval Postgraduate School
ATIN: Mr. G. H. Gleissner	Monterey, CA 93940 1
Bethesda, MD 20084	
Commandant of the Marine Corps	Technical Information Division
(Code AX)	Naval Research Laboratory Washington, D.C. 20375
ATIN: Dr. A. L. Slafkosky	Washington, D.C. 20375
Scientific Advisor	OAST (141), Pentagon
Washington, D.C. 20380 1	ATIN: Mr. Charles S. Smith
	Washington, D.C. 20301 1
Director	washington, D.C. 20301
AMSAA	Reliability Analysis Center
ATIN: DRXSY-MP, H. Cohen	RADC/RBRAC
Aberdeen Proving Ground, MD	ATIN: I. L. Krulac
21005	Data Coordinator/
	Government Programs
	Griffiss AFE. NY 13441 1

Copie	es .	Copies
Dr. Gerhard Heiche Naval Air Systems Command (NAIR 03)	Technical Library Naval Ordinance Station Indian Head, MD 20640	1
Jefferson Plaza No. 1		
Arlington, VA 20360 1 Dr. Barbara Bailar	Library Naval Ocean Systems Center San Diego, CA 92152	1
Associate Director		
Statistical Standards	Technical Library	
Bureau of Census	Bureau of Naval Personnel	
Washington, D.C. 20233 1	Department of the Navy Washington, D.C. 20370	1
Leon Slavin		
Naval Sea Systems Command (NSEA 05H)	Mr. Dan Leonard Code 8105	
Crystal Mall #4, Rm. 129	Naval Ocean Systems Center	_
Washington, D.C. 20036 1	San Diego, CA 92152	1
B. E. Clark RR #2, Box 647-B	Dr. Alan F. Petty Code 7930	
Graham, NC 27253 1	Naval Research Laboratory Washington, D.C. 20375	1
Naval Underwater Systems Cente		
ATTN: Dr. Derrill J. Bordelor Code 601		
Newport, RI 02840 1	Fleet Material Support Offi U.S. Navy Supply Center	ce
Naval Electronic Systems Commo (NELEX 612)	and Mechanicsburg, PA 17055	<u> </u>
ATIN: John Schuster	Mr. Ted Tupper	
National Center No. 1	Code M-311C	
Arlington, VA 20360 1	Military Sealift Command	
-	Department of the Navy	_
Defense Logistics Studies Information Exchange	Washington, D.C. 20390	1
Army Logistics Management Cent ATTN: Mr. J. Dowling	ter Dr. Don Gingras Code 733	
Fort Lee, VA 23801 1		1